



TITLE:

N-Fractional Calculus of Some Irrational Functions (Study on Non-Analytic and Univalent Functions and Applications)

AUTHOR(S):

Nishimoto, Katsuyuki

CITATION:

Nishimoto, Katsuyuki. N-Fractional Calculus of Some Irrational Functions (Study on Non-Analytic and Univalent Functions and Applications). 数理解析研究所講究録 2009, 1626: 56-67

ISSUE DATE:

2009-01

URL:

<http://hdl.handle.net/2433/140320>

RIGHT:

N- Fractional Calculus of Some Irrational Functions.

Katsuyuki Nishimoto

Institute for Applied Mathematics, Descartes Press Co.

2 - 13 - 10 Kaguike, Koriyama, 963 - 8833, JAPAN.

Keywords ; Fractional Calculus, N- Fractional Calculus (NFC),
NFC of Products, NFC of Composite Functions

Abstract

In this article N- fractional calculus of the functions (which have multiple root signs)

$$f(z) = \sqrt{\sqrt{\sqrt{z-b}-c}-d}$$

are discussed .

Theorem 1. We have

$$(i) \quad (f(z))_\gamma = e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k \Gamma(\frac{k}{2}-\frac{1}{8}+\frac{m}{4}+\gamma)}{m! \cdot k! \Gamma(\frac{k}{2}-\frac{1}{8}+\frac{m}{4})} \\ \times \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m, \quad \left(\left| \frac{\Gamma(\frac{k}{2}-\frac{1}{8}+\frac{m}{4}+\gamma)}{\Gamma(\frac{k}{2}-\frac{1}{8}+\frac{m}{4})} \right| < \infty \right)$$

and

$$(ii) \quad (f(z))_n = (-1)^n (z-b)^{(1/8)-n} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k [\frac{k}{2}-\frac{1}{8}+\frac{m}{4}]_n}{m! \cdot k!} \\ \times \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m, \quad (n \in \mathbb{Z}_0^+)$$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1, \quad \left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

§ 0. Introduction (Definition of Fractional Calculus)

(I) Definition. (by K. Nishimoto) ([1] Vol. 1)

Let $D = \{D_-, D_+\}$, $C = \{C_-, C_+\}$,

C_- be a curve along the cut joining two points z and $-\infty + i\text{Im}(z)$,

C_+ be a curve along the cut joining two points z and $\infty + i\text{Im}(z)$,

D_- be a domain surrounded by C_- , D_+ be a domain surrounded by C_+ .

(Here D contains the points over the curve C).

Moreover, let $f = f(z)$ be a regular function in $D(z \in D)$,

$$f_\nu = (f)_\nu = {}_C(f)_\nu = \frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{f(\xi)}{(\xi-z)^{\nu+1}} d\xi \quad (\nu \notin \mathbb{Z}), \quad (1)$$

$$(f)_{-m} = \lim_{\nu \rightarrow -m} (f)_\nu \quad (m \in \mathbb{Z}^+), \quad (2)$$

where $-\pi \leq \arg(\xi-z) \leq \pi$ for C_- , $0 \leq \arg(\xi-z) \leq 2\pi$ for C_+ ,

$\xi \neq z$, $z \in C$, $\nu \in \mathbb{R}$, Γ ; Gamma function,

then $(f)_\nu$ is the fractional differintegration of arbitrary order ν (derivatives of order ν for $\nu > 0$, and integrals of order $-\nu$ for $\nu < 0$), with respect to z , of the function f , if $|(f)_\nu| < \infty$.

(II) On the fractional calculus operator N^ν [3]

Theorem A. Let fractional calculus operator (Nishimoto's Operator) N^ν be

$$N^\nu = \left(\frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{d\xi}{(\xi-z)^{\nu+1}} \right) \quad (\nu \notin \mathbb{Z}), \quad [\text{Refer to (1)}] \quad (3)$$

with
$$N^{-m} = \lim_{\nu \rightarrow -m} N^\nu \quad (m \in \mathbb{Z}^+), \quad (4)$$

and define the binary operation \circ as

$$N^\beta \circ N^\alpha f = N^\beta N^\alpha f = N^\beta (N^\alpha f) \quad (\alpha, \beta \in \mathbb{R}), \quad (5)$$

then the set

$$\{N^\nu\} = \{N^\nu | \nu \in \mathbb{R}\} \quad (6)$$

is an Abelian product group (having continuous index ν) which has the inverse transform operator $(N^\nu)^{-1} = N^{-\nu}$ to the fractional calculus operator N^ν , for the function f such that $f \in F = \{f; 0 \neq |f_\nu| < \infty, \nu \in \mathbb{R}\}$, where $f = f(z)$ and $z \in C$.
(vis. $-\infty < \nu < \infty$).

(For our convenience, we call $N^\beta \circ N^\alpha$ as product of N^β and N^α .)

Theorem B. " F.O.G. $\{N^\nu\}$ " is an " Action product group which has continuous index ν " for the set of F . (F.O.G. ; Fractional calculus operator group) [3]

(III) **Lemma.** We have [1]

$$(i) \quad ((z-c)^b)_\alpha = e^{-i\pi\alpha} \frac{\Gamma(\alpha-b)}{\Gamma(-b)} (z-c)^{b-\alpha} \quad \left(\left| \frac{\Gamma(\alpha-b)}{\Gamma(-b)} \right| < \infty \right), \quad (7)$$

$$(ii) \quad (\log(z-c))_\alpha = -e^{-i\pi\alpha} \Gamma(\alpha) (z-c)^{-\alpha} \quad (|\Gamma(\alpha)| < \infty), \quad (8)$$

$$(iii) \quad ((z-c)^{-\alpha})_{-\alpha} = -e^{i\pi\alpha} \frac{1}{\Gamma(\alpha)} \log(z-c) \quad (|\Gamma(\alpha)| < \infty), \quad (9)$$

where $z-c \neq 0$ for (i) and $z-c \neq 0, 1$ for (ii), (iii),

§ 1. Preliminary

[I] The theorem below is reported by K. Nishimoto already (cf. J. Frac. Calc. Vol. 29, May (2006), p.37). [12]

Theorem D. We have

$$(i) \quad (((z-b)^\beta - c)^\alpha)_\gamma = e^{-i\pi\gamma} (z-b)^{\alpha\beta-\gamma}$$

$$\times \sum_{k=0}^{\infty} \frac{[-\alpha]_k \Gamma(\beta k - \alpha\beta + \gamma)}{k! \Gamma(\beta k - \alpha\beta)} \left(\frac{c}{(z-b)^\beta} \right)^k \quad \left(\left| \frac{\Gamma(\beta k - \alpha\beta + \gamma)}{\Gamma(\beta k - \alpha\beta)} \right| < \infty \right) \quad (1)$$

and

$$(ii) \quad (((z-b)^\beta - c)^\alpha)_n = (-1)^n (z-b)^{\alpha\beta-n}$$

$$\times \sum_{k=0}^{\infty} \frac{[-\alpha]_k [\beta k - \alpha\beta]_n}{k!} \left(\frac{c}{(z-b)^\beta} \right)^k \quad (n \in \mathbb{Z}_0^+) \quad (2)$$

where

$$|c/(z-b)^\beta| < 1,$$

and

$$[\lambda]_k = \lambda(\lambda+1) \cdots (\lambda+k-1) = \Gamma(\lambda+k)/\Gamma(\lambda) \quad \text{with} \quad [\lambda]_0 = 1.$$

(Notation of Pochhammer).

[11] The theorem below is reported by K. Nishimoto already (cf. J. Frac. Calc. Vol. 31, May (2007), p.13). [13]

Theorem E. We have

$$\begin{aligned}
 (i) \quad & (((z-b)^\beta - c)^\alpha - d)^\delta)_\gamma = e^{-i\pi\gamma} (z-b)^{\alpha\beta\delta-\gamma} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k \Gamma(\beta k - \alpha\beta(\delta-m) + \gamma)}{m! \cdot k! \Gamma(\beta k - \alpha\beta(\delta-m))} \left(\frac{c}{(z-b)^\beta} \right)^k \left(\frac{d}{(z-b)^{\alpha\beta}} \right)^m \\
 & \left(\left| \frac{\Gamma(\beta k - \alpha\beta(\delta-m) + \gamma)}{\Gamma(\beta k - \alpha\beta(\delta-m))} \right| < \infty \right)
 \end{aligned} \tag{3}$$

and

$$\begin{aligned}
 (ii) \quad & (((z-b)^\beta - c)^\alpha - d)^\delta)_n = (-1)^n (z-b)^{\alpha\beta\delta-n} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k [\beta k - \alpha\beta(\delta-m)]_n}{m! \cdot k!} \left(\frac{c}{(z-b)^\beta} \right)^k \left(\frac{d}{(z-b)^{\alpha\beta}} \right)^m \\
 & (n \in \mathbb{Z}_0^+)
 \end{aligned} \tag{4}$$

where

$$((z-b)^\beta - c)^\alpha - d \neq 0, \quad |c/(z-b)^\beta| < 1, \quad |d/(z-b)^{\alpha\beta}| < 1.$$

§ 2. N- Fractional Calculus of Functions

$$\sqrt{\sqrt{\sqrt{z-b-c-d}}}$$

Theorem 1. We have

$$\begin{aligned}
 (i) \quad & \left(\sqrt{\sqrt{\sqrt{z-b-c-d}}} \right)_\gamma = e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \\
 & \left(\left| \frac{\Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{\Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \right| < \infty \right)
 \end{aligned} \tag{1}$$

and

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

$$\begin{aligned}
 (ii) \quad & \left(\sqrt{\sqrt{\sqrt{z-b-c-d}}} \right)_n = (-1)^n (z-b)^{(1/8)-n} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k [\frac{k}{2}-\frac{1}{8}+\frac{m}{4}]_n}{m! \cdot k!} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (2) \\
 & (n \in \mathbb{Z}_0^+)
 \end{aligned}$$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1, \quad \left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

Proof of (i). We have

$$\sqrt{\sqrt{\sqrt{z-b-c-d}}} = (((z-b)^{1/2} - c)^{1/2} - d)^{1/2}, \quad (3)$$

hence, operating N^γ to the both sides of (3), we obtain

$$\left(\sqrt{\sqrt{\sqrt{z-b-c-d}}} \right)_\gamma = \left((((z-b)^{1/2} - c)^{1/2} - d)^{1/2} \right)_\gamma \quad (4)$$

$$= e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k \Gamma(\frac{k}{2}-\frac{1}{8}+\frac{m}{4}+\gamma)}{m! \cdot k! \Gamma(\frac{k}{2}-\frac{1}{8}+\frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (5)$$

setting $\alpha = \beta = \delta = 1/2$ in Theorem E. (i) in preliminary, under the conditions stated before.

Proof of (ii). Set $\gamma = n$ in (i).

Corollary 1. We have

$$\begin{aligned}
 (i) \quad & \left(\sqrt{\sqrt{\sqrt{z-b-c}}} \right)_\gamma = e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma} \\
 & \times \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k \Gamma(\frac{k}{2}-\frac{1}{8}+\gamma)}{k! \Gamma(\frac{k}{2}-\frac{1}{8})} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (6) \\
 & \left(\left| \frac{\Gamma(\frac{k}{2}-\frac{1}{8}+\gamma)}{\Gamma(\frac{k}{2}-\frac{1}{8})} \right| < \infty \right)
 \end{aligned}$$

and

$$(ii) \quad \left(\sqrt{\sqrt{\sqrt{z-b}-c}} \right)_n = (-1)^n (z-b)^{(1/8)-n} \times \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k [\frac{k}{2}-\frac{1}{8}]_n}{k!} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (7)$$

$(n \in \mathbb{Z}_0^+)$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1.$$

Proof of (i). Set $d = 0$ in Theorem 1 (i).

We have (6) from Theorem D, (i), setting $\beta = 1/2$ and $\alpha = 1/4$.

Proof of (ii). Set $\gamma = n$ in (i).

Corollary 2. We have

$$(i) \quad \left(\sqrt{\sqrt{\sqrt{z-b}}} \right)_\gamma = e^{-i\pi\gamma} \frac{\Gamma(-\frac{1}{8}+\gamma)}{\Gamma(-\frac{1}{8})} (z-b)^{(1/8)-\gamma} \quad (8)$$

$(|\Gamma(-\frac{1}{8}+\gamma)| < \infty)$

and

$$(ii) \quad \left(\sqrt{\sqrt{\sqrt{z-b}}} \right)_n = (-1)^n [-\frac{1}{8}]_n (z-b)^{(1/8)-n} \quad (9)$$

$(n \in \mathbb{Z}_0^+)$

where

$$z-b \neq 0.$$

Proof of (i). Set $c = 0$ in Corollary 1 (i).

Or set $c = d = 0$ in Theorem 1. (i).

Proof of (ii). Set $\gamma = n$ in (i).

Note. We have

$$\left(\sqrt{\sqrt{\sqrt{z-b}}} \right)_\gamma = ((z-b)^{1/8})_\gamma = e^{-i\pi\gamma} \frac{\Gamma(-\frac{1}{8}+\gamma)}{\Gamma(-\frac{1}{8})} (z-b)^{(1/8)-\gamma} \quad (10)$$

from Lemma (i), directly.

§ 3. Semi Derivatives and Integrals

Theorem 2. We have

$$(i) \quad \left(\sqrt{\sqrt{\sqrt{z-b-c-d}}} \right)_{1/2} = -i(z-b)^{-3/8} \\ \times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} + \frac{3}{8} + \frac{m}{4})}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (1) \\ \text{(semi derivatives)}$$

and

$$(ii) \quad \left(\sqrt{\sqrt{\sqrt{z-b-c-d}}} \right)_{-1/2} = i(z-b)^{5/8} \\ \times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{5}{8} + \frac{m}{4})}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (2) \\ \text{(semi integrals)}$$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1, \quad \left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

Proof. Set $\gamma = 1/2$ and $-1/2$ in Theorem 1(i) we have then (i) and (ii) respectively, clearly.

Corollary 3. We have

$$(i) \quad \left(\sqrt{\sqrt{\sqrt{z-b-c}}} \right)_{1/2} = -i(z-b)^{-3/8} \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k \Gamma(\frac{k}{2} + \frac{3}{8})}{k! \Gamma(\frac{k}{2} - \frac{1}{8})} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (3) \\ \text{(semi derivatives)}$$

and

$$(ii) \quad \left(\sqrt{\sqrt{\sqrt{z-b-c}}} \right)_{-1/2} = i(z-b)^{5/8} \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{5}{8})}{k! \Gamma(\frac{k}{2} - \frac{1}{8})} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (4) \\ \text{(semi integrals)}$$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1.$$

Proof. Set $d = 0$ in Theorem 2.

Corollary 4. We have

$$(i) \quad \left(\sqrt{\sqrt{\sqrt{z-b}} - d} \right)_{1/2} = -i(z-b)^{-3/8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m \Gamma(\frac{3}{8} + \frac{m}{4})}{m!! \Gamma(-\frac{1}{8} + \frac{m}{4})} \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (5)$$

(semi derivatives)

and

$$(ii) \quad \left(\sqrt{\sqrt{\sqrt{z-b}} - d} \right)_{-1/2} = i(z-b)^{5/8} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m \Gamma(-\frac{5}{8} + \frac{m}{4})}{m! \Gamma(-\frac{1}{8} + \frac{m}{4})} \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (2)$$

(semi integrals)

where

$$\left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

Proof. Set $c = 0$ in Theorem 2.

§ 4. Some Special Cases

[I] When $n = 0$, we have

$$\begin{aligned} \left(\sqrt{\sqrt{\sqrt{z-b-c-d}}} \right)_0 &= (z-b)^{1/8} \\ &\times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k}{m! \cdot k!} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \end{aligned} \quad (1)$$

from § 2. (2).

Now we have

$$\text{RHS of (1)} = (z-b)^{1/8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k}{k!} S^k \quad (2)$$

$$\left(S = \frac{c}{\sqrt{z-b}}, \quad T = \frac{d}{\sqrt[4]{z-b}} \right)$$

$$= (z-b)^{1/8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m (1-S)^{(1/4)-(m/2)} \quad (3)$$

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

$$= (z-b)^{1/8} (1-S)^{1/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m T^m}{m! (1-S)^{m/2}} \quad (4)$$

$$= (z-b)^{1/8} (1-S)^{1/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} \left(\frac{d}{\sqrt{\sqrt{z-b}-c}} \right)^m \quad (5)$$

$$= (z-b)^{1/8} \left(1 - \frac{c}{\sqrt{z-b}} \right)^{1/4} \left(1 - \frac{d}{\sqrt{\sqrt{z-b}-c}} \right)^{1/2} \quad (6)$$

$$= \sqrt{\sqrt{\sqrt{z-b}-c}-d} \quad (\text{LHS of (1)}) \quad (7)$$

[II] When $n=1$, we have

$$\left(\sqrt{\sqrt{\sqrt{z-b}-c}-d} \right)_1 = -(z-b)^{-7/8} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k [\frac{k}{2}+\frac{m}{4}-\frac{1}{8}]_1}{m! \cdot k!} S^k T^m \quad (8)$$

$$= -(z-b)^{-7/8} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k (\frac{k}{2}+\frac{m}{4}-\frac{1}{8})}{m! \cdot k!} S^k T^m \quad (9)$$

from § 2. (2).

Now we have the identities;

$$\sum_{k=0}^{\infty} \frac{[\lambda]_k}{k!} z^k = (1-z)^{-\lambda}, \quad (10)$$

$$[\lambda]_0 = 1, \quad [\lambda]_1 = \lambda, \quad (11)$$

$$[\lambda]_{k+1} = \lambda [\lambda+1]_k, \quad (12)$$

hence

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{[\lambda]_k k}{k!} z^k &= \sum_{k=1}^{\infty} \frac{[\lambda]_k}{(k-1)!} z^k = z \sum_{k=0}^{\infty} \frac{[\lambda]_{k+1}}{k!} z^k \\ &= \lambda z (1-z)^{-\lambda-1}. \end{aligned} \quad (13)$$

Therefore, we have

$$\sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2}-\frac{1}{4}]_k (\frac{k}{2})}{m! \cdot k!} S^k T^m = \frac{1}{2} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2}-\frac{1}{4}]_k k}{k!} S^k \quad (14)$$

$$= \frac{1}{2} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \left(\frac{m}{2} - \frac{1}{4} \right) S \sum_{k=0}^{\infty} \frac{[\frac{m}{2} + \frac{3}{4}]_k}{k!} S^k \quad (15)$$

$$= \frac{1}{2} S(1-S)^{-3/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m (\frac{m}{2} - \frac{1}{4})}{m!} U^m \quad \left(U = \frac{d}{\sqrt{\sqrt{z-b}-c}} \right) \quad (16)$$

$$= \frac{1}{2} S(1-S)^{-3/4} \left\{ -\frac{1}{4} U(1-U)^{-1/2} - \frac{1}{4} (1-U)^{1/2} \right\} \quad (17)$$

$$= -\frac{1}{8} S(1-S)^{-3/4} (1-U)^{-1/2}, \quad (18)$$

$$\sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k (\frac{m}{4})}{m! k!} S^k T^m = \frac{1}{4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k}{k!} S^k \quad (19)$$

$$= \frac{1}{4} (1-S)^{1/4} \sum_{m=1}^{\infty} \frac{[-\frac{1}{2}]_m}{(m-1)!} U^m \quad (20)$$

$$= \frac{1}{4} (1-S)^{1/4} U \left(-\frac{1}{2} \right) \sum_{m=0}^{\infty} \frac{[\frac{1}{2}]_m}{m!} U^m \quad (21)$$

$$= -\frac{1}{8} (1-S)^{1/4} U(1-U)^{-1/2}. \quad (22)$$

and

$$\sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k (-\frac{1}{8})}{m! k!} S^k T^m = -\frac{1}{8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k}{k!} S^k \quad (23)$$

$$= -\frac{1}{8} (1-S)^{1/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} U^m \quad (24)$$

$$= -\frac{1}{8} (1-S)^{1/4} (1-U)^{1/2}. \quad (25)$$

Then applying (18), (22) and (25) to (9) we obtain

$$\left(\sqrt{\sqrt{\sqrt{z-b}-c}-d} \right)_1 = \frac{1}{8} (z-b)^{-7/8} (1-S)^{1/4} (1-U)^{-1/2} \{ S(1-S)^{-1} + 1 \} \quad (26)$$

$$= \frac{1}{8}(z-b)^{-7/8}(1-S)^{1/4}(1-U)^{-1/2}(1-S)^{-1} \quad (27)$$

$$= \frac{1}{8}(z-b)^{-7/8}(1-S)^{-3/4}(1-U)^{-1/2} \quad (28)$$

$$= \frac{1}{8}(z-b)^{-1/2}(\sqrt{z-b}-c)^{-1/2}(\sqrt{\sqrt{z-b}-c}-d)^{-1/2} \quad (29)$$

This result (29) coincides with the one obtained by classical calculus.

References

- [1] K. Nishimoto ; Fractional Calculus, Vol. 1 (1984), Vol. 2 (1987), Vol. 3 (1989), Vol. 4 (1991), Vol. 5, (1996), Descartes Press, Koriyama, Japan.
- [2] K. Nishimoto ; An Essence of Nishimoto's Fractional Calculus (Calculus of the 21st Century); Integrals and Differentiations of Arbitrary Order (1991), Descartes Press, Koriyama, Japan.
- [3] K. Nishimoto ; On Nishimoto's fractional calculus operator N^\vee (On an action group), J. Frac. Calc. Vol. 4, Nov. (1993), 1 - 11.
- [4] K. Nishimoto ; Unification of the integrals and derivatives (A serendipity in fractional calculus), J. Frac. Calc. Vol. 6, Nov. (1994), 1 - 14.
- [5] K. Nishimoto ; Ring and Field Produced from The Set of N- Fractional Calculus Operator, J. Frac Calc. Vol. 24, Nov. (2003), 29 - 36.
- [6] K. Nishimoto ; On the fractional calculus $(a-z)^b$ and $\log(a-z)$, J. Frac. Calc. Vol.3, May (1993), 19 - 27.
- [7] K. Nishimoto and S.- T. Tu ; Fractional calculus of Psi functions (Generalized Polygamma unctions), J. Frac. Calc. Vol.5 May (1994), 27 - 34.
- [8] S.- T. Tu and K. Nishimoto ; On the fractional calculus of functions $(cz-a)^b \log(cz-a)$, J. Frac.Calc.Vol.5, May (1994), 35 - 43.
- [9] K. Nishimoto ; N- Fractional Calculus of the Power and Logarithmic Functions and Some Identities, J. Frac. Calc. Vol.21, May (2002), 1 - 6.
- [10] K. Nishimoto ; Some Theorems for N- Fractional Calculus of Logarithmic Functions I, J. Frac Calc.Vol.21, May (2002), 7 - 12.
- [11] K. Nishimoto ; N- Fractional Calculus of Products of Some Power Functions, J. Frac.Calc. Vol.27, May (2005), 83 - 88.
- [12] K. Nishimoto ; N- Fractional Calculus of Some Composite Functions, J. Frac. Calc. Vol. 29, May (2006), 35 - 44.
- [13] K. Nishimoto ; N- Fractional Calculus of Some Composite Algebraic Functions, J. Frac.Calc. Vol. 31, May (2006), 11 - 23.
- [14] K. Nishimoto ; N- Fractional Calculus of Some Elementary Functions and Their Semi Differintegrations, J. Frac. Caic. Vol. 31, May (2007), 1 - 10.
- [15] K. Nishimoto and T. Miyakoda ; N- Fractional Calculus and n -th Derivatives of Some Algebraic Functions, J. Frac. Calc. Vol. 31, May (2007), 53 - 62.

Katsuyuki Nishimoto

- [16] T. Miyakoda ; N-Fractional Calculus of Certain Algebraic Functions, J. Frac. Calc. Vol. 31, May (2007), 63 - 76.
- [17] David Dunmmit and Richard M. Foote ; Abstract Algebra, Prentice Hall (1991).
- [18] K. B. Oldham and J. Spanier ; The Fractional Calculus, Academic Press (1974).
- [19] A.C. McBride ; Fractional Calculus and Integral Transforms of Generalized Functions, Research Notes, Vol. 31, (1979), Pitman.
- [20] S.G. Samko, A.A. Kilbas and O.I. Marichev ; Fractional Integrals and Derivatives, and Some Their Applications (1987), Naoka, USSR.
- [21] K.S. Miller and B. Ross ; An Introduction to The Fractional Calculus and Fractional Diffetial Equations, John Wiley & Sons, (1993).
- [22] V. Kiryakova ; Generalized fractional calculus and applications, Pitman Research Notes, No.301, (1994), Longman.
- [23] A.Carpinteri and F. Mainardi (Ed.) ; Fractals and Fractional Calculus in Continume Mechanics (1997), Springer, Wien, New York.
- [24] Igor Podlubny ; Fractional Differential Equations (1999), Academic Press.
- [25] R. Hilfer (Ed.) ; Applications of Fractional Calculus in Physics, (2000), World Scientific, Singapor, New Jersey, London, Hong Kong.
- [26] A.P. Prudnikov, Yu. A. Bryckov and O.I. Marichev ; Integrals and Series, Vol. I, Gordon and Breach, New York, (1986).
- [27] S. Moriguchi, K.Udagawa and S. Hitotsumatsu ; Mathematical Formulae, Vol.2, Iwanami Zensho, (1957), Iwanami, Japan.

Katsuyuki Nishimoto
 Institute for Applied Mathematics
 Descartes Press Co.
 2 - 13 - 10 Kaguike, Koriyama
 963 - 8833 Japan